

Supplemental Approvals

Generic Name (Trade Name) Company

October 29, 2018

Sodium oxybate

Uses/Notes

[FDA approved sodium oxybate](#) for treatment of cataplexy and excessive daytime sleepiness in pediatric patients (aged 7–17 y) with narcolepsy.

Sodium oxybate is a central nervous system (CNS) depressant approved in 2002 for treatment of cataplexy in adult patients with narcolepsy. Cataplexy is a sudden and transient episode of muscle weakness accompanied by full conscious awareness, typically triggered by emotions such as laughing, crying, or terror.

Sodium oxybate either alone or in combination with other CNS depressants may be associated with adverse reactions that include seizure, respiratory depression, decreases in the level of consciousness, coma, and death. Rapid onset of sedation, coupled with amnesia, particularly when combined with alcohol, has posed risks for voluntary and involuntary users (e.g., assault victims).

The agent is contraindicated in patients being treated with sedative hypnotic agents and in patients with succinic semialdehyde dehydrogenase deficiency. In addition, patients should not drink alcohol when using Sodium oxybate. Succinic semialdehyde deficiency is a rare inborn error of metabolism variably characterized by mental retardation, hypotonia, and ataxia.

The most common adverse reactions in pediatric patients were bed-wetting, nausea, headache, vomiting, weight decrease, decreased appetite, and dizziness.

The following adverse reactions have been identified during postapproval use of sodium oxybate. Because these reactions are reported voluntarily from a population of uncertain size, it is not always possible to reliably estimate their frequency or establish a causal relationship to drug exposure: joint pain, decreased appetite, fall, fluid retention, hangover, headache, hypersensitivity, hypertension, memory impairment, nocturia, panic attack, vision blurred and weight decreased.

(Xyrem—Jazz Pharmaceuticals)

Agent now approved for treatment of cataplexy and excessive daytime sleepiness in children

Because of the risk of serious outcomes resulting from inappropriate prescribing, misuse, abuse and diversion, sodium oxybate is only available through a [risk evaluation mitigation strategy \(REMS\) programs](#).

Source URL:

<http://aphanet.org/supplemental-approvals/agent-now-approved-treatment-cataplexy-and-excessive-daytime-sleepiness>

APhA DrugInfoLine is an official publication of, and is owned and copyrighted by the American Pharmacists Association, the national professional society of pharmacists. Materials in APhA DrugInfoLine do not necessarily represent the policy, recommendations, or endorsement of APhA. The publisher, authors, editors, reviewers, and contributors have taken care to ensure that information contained in APhA DrugInfoLine is accurate and current; however, they shall have no liability to any person or entity with regard to claims, losses, or damages caused or alleged to be caused, directly or indirectly, by use of any information contained in the publication. All decisions about drug therapy must be based on the independent judgment of the clinician. Copyright © 2000–2011, American Pharmacists Association. All rights reserved.